



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GENERAL NOTES.

The Black Tern (*Hydrochelidon nigra surinamensis*) in New York Harbor.—On August 31, 1899, five Black Terns were shot at Bedloes Island, New York Harbor. The two that I skinned, both males, had been eating small fish. Others of the same species were seen subsequently in the same locality at about same date. — L. S. FOSTER, *New York City*.

Capture of the Brown Pelican in Wyoming.—I have to record the capture of *Pelecanus fuscus* upon one of the city reservoirs of Cheyenne, Wyoming, on July 12, one individual, a female, which was alone. The bird was in poor flesh and apparently had about completed moulting. Two or three of her rectrices were less than half grown and each wing showed two or three old brown primaries and two or three old brown secondaries not yet shed. Head and neck uniform slate gray. Measurements as follows: Culmen, 11.12; wing, 20.00; length, 49.00 inches. I do not know of any other captures of this species in Wyoming, or, for that matter, in any intermountain State, in this or contiguous latitudes. I am not familiar with the species, but think the uniform brown head and neck would indicate immaturity, but the few worn and much darker quills of the wings and the partly moulted tail offer conclusive evidence against the suggestion that the bird is of the present year. I have mounted the specimen and placed it in the Cheyenne High School collection. — FRANK BOND, *Cheyenne, Wyo.*

The Little Blue Heron (*Ardea cærulea*) in Connecticut.—On July 21, 1899, I saw no less than seven individuals of this species, all young birds, in white plumage. On the 25th I secured one, and a few days later another; there were five shot all together. The remaining two left and were not seen again. The flock spent about two weeks here, feeding on the salt marshes. This is the first and only time that I have met with *Ardea cærulea*, in Connecticut. — ARTHUR WILLIAM BROCKWAY, *Old Lyme, Conn.*

The Green Heron Breeding in Ontario.—Very little seems to be known of the breeding habits of the Green Heron (*Ardea virescens*) in Ontario, and its presence is not often noted north of the southern parts of the Province—McIlwraith makes the general statement that it breeds near Dunville, the St. Clair Flats, and at Hamilton. This summer we have located it in a spot fully eighty miles north of these places and there is reason to believe that it breeds at several points in this locality. Early in June the Rev. C. J. Young of Lansdowne, Ont., informed me